

“Routine circumcision is not a medical issue or a social issue. It is a sexual issue and a human rights issue.”
Frederick Hodges

How is circumcision done?

Most parents don’t know what is actually done to a baby when he is circumcised. The baby is placed spread-eagle on his back on a board and his arms and legs are strapped down so that he can’t move. His genitals are scrubbed and covered with antiseptic. His foreskin is torn from his glans and slit lengthwise so that the circumcision instrument can be inserted. Then his foreskin is cut off.

Most parents who see what is done to a baby when he is circumcised and how he reacts decide against circumcision and let their baby keep his foreskin intact.

Parents have new concerns

More and more parents – including Jewish parents – are questioning the wisdom of subjecting their baby to the pain and risks of circumcision and its life-long consequences. More and more parents are wondering if they have the right to consent to the irreversible amputation of a healthy, normal, sensitive, functional part of their baby’s penis – an amputation that experts regard not just as unnecessary, but as contra-indicated. More and more parents are becoming truly informed and, as a result, more and more parents are deciding against circumcision and are keeping their baby boys intact.

“The best reason to let a baby keep his foreskin intact is that it’s almost a certainty he will be glad you did.”
John A. Erickson

“Many parents today realize that if they had been given accurate information about circumcision, they would never have let anyone circumcise their baby. I am one of those parents, and that is why I do the work I do and why I have written this pamphlet.”
Marilyn Fayre Milos, R.N.

More information can be found on the Internet at: <http://www.nocirc.org/> and <http://www.cirp.org/CIRP/>.

NOCIRC pamphlets: 50¢ each or \$25/100 (same or mixed) plus \$3 S/H. Five different pamphlets: \$2 and business-size SASE with 55¢ postage.

The **NOCIRC Resource Guide** lists the pamphlets, books, articles, newsletters and videos available from NOCIRC, and other resources as well. Included as one of the five pamphlets above or free for SASE.

National Organization of Circumcision Information Resource Centers

Post Office Box 2512
San Anselmo, CA 94979-2512 USA

Telephone: 415-488-9883
Fax: 415-488-9660

www.nocirc.org



The information in this pamphlet is not meant to replace the care and advice of your pediatrician.

Answers To Your Questions About Infant Circumcision

from the
**National Organization
of Circumcision Information
Resource Centers**

*Educating a New Generation
For the Well-Being of All Children*

www.nocirc.org

What is circumcision?

Circumcision is the cutting off of the fold of skin that normally covers the glans of the penis. This double layer of skin, the prepuce, is commonly known as the foreskin.

Why is the foreskin there?

The foreskin comprises as much as half or more of the penile skin system and has three known functions: protective, sensory and sexual.

During infancy, the foreskin is attached to the glans and protects it from urine, feces and abrasion from diapers. Throughout life, the foreskin keeps the glans soft and moist and protects it from trauma and injury. Without this protection, the glans becomes dry, calloused and desensitized from exposure and chafing.

Specialized nerve endings in the foreskin enhance sexual pleasure.

The foreskin may have functions not yet recognized or understood.

When and why did doctors in the U.S. start circumcising babies?

Doctors in the English-speaking countries started circumcising babies in the mid-1800s “to prevent masturbation,” which was blamed for causing many diseases, including epilepsy, tuberculosis and insanity.

Other reasons have been given since then, but all of them, including the claim that circumcision prevents cancer of the penis, cancer of the cervix, and venereal diseases, have been disproven. We now know that the foreskin is a normal, sensitive, functional part of the body.

If my son isn't circumcised, won't he feel different? Won't he be teased?

With more and more males growing up intact, it's circumcised males who are beginning to feel different. Many circumcised men have reported that they have been teased because they didn't have a foreskin.

Is circumcision painful?

Yes. Circumcision is extremely painful – and traumatic – for a baby. Just being strapped down is frightening for a baby. The often repeated statement that babies can't feel pain is not true. Babies are as sensitive to pain as anyone else. Most babies scream frantically when their foreskins are cut off. Some defecate. Some lapse into a coma. The reason some babies don't cry when they are circumcised is that they *can't* cry because they are in a state of shock. Most babies are circumcised without an anesthetic. Anesthetics injected into the penis don't always work. Being stuck with a needle in the penis is itself painful for a baby, just as it would be for anyone else. Babies are rarely given pain medication right after they are circumcised or during the week to ten days it takes the wound to heal. Pain medication is not always effective and is never 100% effective.

Does circumcision have risks?

Yes. Like any other surgery, circumcision has risks. They include:

- Excessive bleeding
- Infection
- Complications from anesthetics
- Surgical mistakes, including loss of glans and loss of entire penis
- Death

Many circumcised males suffer from:

- Extensive scarring
- Skin tags and skin bridges
- Tearing and bleeding at the scar
- Curvature of the penis
- Tight, painful erections
- Difficulty ejaculating
- Impotence
- Feelings of having been violated
- Feelings of having been mutilated

All circumcised males lose some or most of the sensitivity in their glans and *all* of the sensitivity in their foreskins.

Circumcision may have risks and complications not yet recognized or understood.